

On topological properties of compact attractors on Hausdorff spaces

Wouter Jongeneel

Abstract—We characterize when a compact, invariant, asymptotically stable attractor on a locally compact Hausdorff space is a strong deformation retract of its domain of attraction.

I. INTRODUCTION

The purpose of this note is to improve our understanding of topological properties of compact asymptotically stable attractors and their respective domain of attraction. Here, we will almost exclusively appeal to topological tools pioneered by Borsuk [1]. In particular, we will elaborate on the *retraction theoretic* work by Moulay & Bhat [2], which itself is a generalization of the seminal works [3, Thm. 21], [4] and [5, Thm. 4.1].

After Poincaré and Lyapunov (Liapunov), the modern qualitative study of attractors was largely propelled through the monographs by Birkhoff [6] and Nemytskii & Stepanov [7], with influential follow-up works by Auslander, Bhatia & Siebert [8], Wilson [9], Hahn [10], Bhatia & Szegő [11], Conley [12], Milnor [13] and many others, *e.g.*, see [14, Ch. 1].

Lately, attractors have been extensively studied through the lens of *shape theory*, *e.g.*, see [15]–[17] and [18, Prop. 1], with the seminal work of Günther & Segal showing that a finite-dimensional compact subset of a manifold can be an attractor if and only if it has the shape of a finite polyhedron [19].

The interest in understanding topological properties of attractors and their respective domain of attraction stems from the simple observation that if a certain dynamical system does not exist, then certainly there is no feedback law resulting in a closed-loop dynamical system with precisely those dynamics.

Indeed, this type of study often provides *necessary* conditions of the form that an attractor must be *equivalent* in some sense to its domain of attraction. With that in mind, one seeks a notion of equivalence that is weak enough to cover many dynamical systems, yet also strong enough to obtain insights, *e.g.*, obstructions. Hence, although shape equivalence is more widely applicable [20] and in that sense more fundamental, we focus on *homotopy equivalence* with the aim of recovering stronger necessary conditions.

In the same spirit, by further restricting the problem class, one could even look for stronger notions of equivalence as recently done in [21]. There, in the context of a vector-field

guided path-following problem, homotopy equivalences have been strengthened to topological equivalences (homeomorphisms).

Although we focus on continuous dynamical systems, one can link work of this form to families of differential inclusions [22]. Indeed, further partial generalizations of [2] to nonsmooth dynamical system are presented in [23].

Notation and technical preliminaries: The *identity map* on a space X is denoted by id_X , that is, $\text{id}_X : x \mapsto x \ \forall x \in X$. The (embedded) n -sphere is the set $\mathbb{S}^n := \{x \in \mathbb{R}^{n+1} : \langle x, x \rangle = 1\}$, with the (closed) n -disk being $\mathbb{D}^n := \{x \in \mathbb{R}^n : \langle x, x \rangle \leq 1\}$. The topological boundary of a space X is denoted by ∂X , *e.g.*, $\partial \mathbb{D}^{n+1} = \mathbb{S}^n$. We use \simeq_h to denote homotopy (equivalence), see Section. II-B.

A topological space X is said to be a *locally compact Hausdorff* space when: (i) for any $x \in X$ there is a compact set $K \subseteq X$ containing an open neighbourhood U of x and; (ii) for any $x_1, x_2 \in X$ there are open neighbourhoods $U_1 \ni x_1$ and $U_2 \ni x_2$ such that $U_1 \cap U_2 = \emptyset$, *e.g.*, see [24, p. 31, 104]. Examples of locally compact Hausdorff spaces are: \mathbb{R}^n , topological manifolds, the Hilbert cube, any discrete space and so forth. In particular, any compact Hausdorff space is locally compact. Regarding counterexamples, a space X with the trivial topology $\tau = \{X, \emptyset\}$ is not Hausdorff and any infinite-dimensional Hilbert space is a Hausdorff topological vector space, yet, it fails to be locally compact, see also [25, Thm. 29.1].

II. CONTINUOUS DYNAMICAL SYSTEMS

In this note we study continuous (global) *semi-dynamical systems* comprised of the triple $\Sigma := (M, \varphi, \mathbb{R}_{\geq 0})$. Here, M is a locally compact Hausdorff space and $\varphi : \mathbb{R}_{\geq 0} \times M \rightarrow M$ is a (global) semi-flow, that is, a continuous map that satisfies for any $x \in M$:

- (i) $\varphi(0, x) = x$ (*identity axiom*); and
- (ii) $(\varphi(s, \varphi(t, x))) = \varphi(t + s, x) \ \forall s, t \in \mathbb{R}_{\geq 0} := \{t \in \mathbb{R} : t \geq 0\}$ (*semi-group axiom*).

We will usually write φ^t instead of $\varphi(t, \cdot)$.

We say that a point $x \in M$ is a *start point* (under Σ) if $\forall (t, y) \in \mathbb{R}_{>0} \times M$ we have that $\varphi^t(y) \neq x$. Differently put, $x \in M$ is a starting point when a flow starting from x cannot be extended backwards, see [26, Ex. 5.14] for an example. To avoid confusion, the evaluation of an integral curve at 0 is sometimes called a “*starting point*” [27, p. 206], which is not what we are talking about here. Then, to eliminate the existence of start points we appeal to [26, Prop. 1.7], for instance, we can consider semi-flows generated by a smooth vector field. Concretely, let $F \in \Gamma^\infty(TM)$ be a smooth vector field on a smooth manifold M . It is well-known that

under these conditions, for each $p \in M$ there is a $\varepsilon > 0$ such that $\gamma : (-\varepsilon, \varepsilon) \rightarrow M$ is an integral curve of F with $\gamma(0) = p$ [27, Prop. 9.2], that is, in terms of the (local) flow $\varphi : (-\varepsilon, \varepsilon) \times M \rightarrow M$ we have

$$\left. \frac{d}{dt} \varphi^t(p) \right|_{t=s} = F(\varphi^s(p)), \quad s \in (-\varepsilon, \varepsilon).$$

Hence, with the previous observation in mind we will assume the following throughout the remainder of the note.

Assumption II.1 (Start points). *The set of start points under the semi-dynamical system Σ is empty.*

A. Stability

We will exclusively focus on a subclass of semi-dynamical systems with practically relevant stability properties.

Definition II.2 (Attractor). *Given some global semi-dynamical system $\Sigma = (M, \varphi, \mathbb{R}_{\geq 0})$, then, a compact set $A \subseteq M$ is said to be an invariant, local asymptotically stable, attractor when*

- (i) $\varphi(\mathbb{R}, A) = A$ (invariance); and
- (ii) *for any open neighbourhood $U_\varepsilon \subseteq M$ of A there is an open neighbourhood $U_\delta \subseteq U_\varepsilon \subseteq M$ of A such that $\varphi(\mathbb{R}_{\geq 0}, U_\delta) \subseteq U_\varepsilon$ (Lyapunov stability), plus, there is an open neighbourhood $W \subseteq M$ of A such that all semi-flows initialized in W converge to A (local attractivity), that is, for any $p \in W$ and any open neighbourhood $V \subseteq M$ of A there is a $T \geq 0$ such that $\varphi^t(p) \in V \forall t \geq T$.*

The combination of Lyapunov stability and local attractivity is referred to as *local asymptotic stability*. When the neighbourhood W in Item (ii) can be chosen to be all of M we speak of *global asymptotic stability*. Local asymptotic stability is also captured by the existence of an open neighbourhood $U \subseteq M$ of A such that $\cup_{t \geq 0} \varphi^t(U) = A$ [28, Lem. 1.6]

One can find several definitions of “attractors” in the literature, see for instance [11, Def. V.1.5], [13] and [20, Sec. 2.2].

Definition II.3 (Domain of attraction). *Let the compact set $A \subseteq M$ be an invariant, local asymptotically stable attractor under the semi-dynamical system $\Sigma = (M, \varphi, \mathbb{R}_{\geq 0})$, then, its domain of attraction is*

$$\mathcal{D}_\Sigma(A) = \{p \in M : \text{for any open neighbourhood } U \subseteq M \text{ of } A \text{ there is a } T \geq 0 \text{ such that } \varphi^t(p) \in U \forall t \geq T\}.$$

Definition II.3 can be equivalently written in terms of convergent subsequences. Topological properties of attractors $A \subseteq M$ and their respective domain of attraction $\mathcal{D}_\Sigma(A) \subseteq M$ are an active topic of study since the 1960s [11], [14].

To elaborate on the introduction, the interest stems from the observation that the (numerical) analysis or synthesis, e.g., via feedback control, of dynamical systems Σ can be involved, while topological properties of the pair $(\mathcal{D}_\Sigma(A), A)$

might be readily available. Here, topological knowledge of $\mathcal{D}_\Sigma(A)$ is frequently used to study if some “desirable” domain of attraction is admissible. For instance, one can show that no point $p \in \mathbb{S}^1$ can be a global asymptotically stable attractor under any $\Sigma = (\mathbb{S}^1, \varphi, \mathbb{R}_{\geq 0})$, e.g., see Theorem II.6 below. The intuition being that for this to be true the circle \mathbb{S}^1 needs to be torn apart, which is obstructed by demanding φ to be continuous, see also [14, Fig. 1.1]. Again, we emphasize that conclusions of this form emerge without involved analysis of any particular system Σ .

B. Retraction theory

The previous example can be understood through \mathbb{S}^1 not being *contractible*, that is, \mathbb{S}^1 is not *homotopy equivalent* to a point p . Formally, two topological spaces X and Y are said to have the same *homotopy type* when they are homotopy equivalent¹, that is, there are continuous maps $f : X \rightarrow Y$ and $g : Y \rightarrow X$ such that $f \circ g \simeq_h \text{id}_Y$ and $g \circ f \simeq_h \text{id}_X$. In some sense, this notion is a more general version of a deformation retract—which we recall below, and is most naturally understood through invariance in differential- [29] and algebraic topology [30]. As alluded to, now, we recall that $A \subseteq X$ is a *retract* of X when there is a map $r : X \rightarrow A$ such that $r \circ \iota_A = \text{id}_A$. The set A is said to be a *deformation retract* of X when A is a retract and additionally $\iota_A \circ r \simeq_h \text{id}_X$, implying that X is homotopy equivalent to A . When, additionally, the homotopy is *stationary relative to A* , we speak of a *strong deformation retract*.

Remark II.4 (Deformation retracts). *The literature does not agree on what a “deformation retract” is. For instance, Hatcher calls strong deformation retracts simply deformation retracts and speaks of “deformation retracts in the weak sense” where we would be speak of simply a deformation retract [30, Ch. 0]. This should be contrasted with for instance the 1965 text of Hu [31, Sec. 1.11].*

Next, a set $A \subseteq X$ is said to be a *weak deformation retract* of X when every open neighbourhood $U \supseteq A$ contains a strong deformation retract $V \supseteq A$ of X .

In this note we will elaborate on the following result due to Moulay & Bhat. In particular, we aim to understand when $\mathcal{D}_\Sigma(A)$ strongly deformation retracts onto A and not just to a subset of a neighbourhood around A .

Theorem II.5 ([2, Thm. 5]). *Suppose that the compact set $A \subseteq M$ is an invariant, local asymptotically stable attractor under the semi-dynamical system $\Sigma = (M, \varphi, \mathbb{R}_{\geq 0})$, then, A is a weak deformation retract of $\mathcal{D}_\Sigma(A)$.*

It is well-known that when M is a smooth manifold and A is an embedded submanifold of M , then, A is a strong neighbourhood deformation retract of M and thus A is homotopic to $\mathcal{D}_\Sigma(A)$ [2, Prop. 10], see also [32]. Our aim is to provide further, especially weaker, conditions for this to be true.

¹More abstractly, homotopies are isomorphisms in the homotopy category of topological spaces.

Indeed, Theorem II.5 can be seen as a generalization of the following well-known result due to Sontag.

Theorem II.6 ([3, Thm. 21]). *Suppose that the point $A := \{p\} \subseteq M$ is an invariant, global asymptotically stable attractor under the semi-dynamical system $\Sigma = (M, \varphi, \mathbb{R}_{\geq 0})$, then, $\mathcal{D}_\Sigma(A)$ is contractible.*

We aim to strengthen this generalization. We do this by appealing to neighbourhood retracts. A set $A \subseteq X$ is a *neighbourhood retract* of X when there is an open neighbourhood $U \subseteq X$ of A such that A is retract of U . This definition extends naturally to (strong) deformation retracts. Now indeed, if for instance M weakly deformation retracts onto $A \subseteq M$ while A is a neighbourhood deformation retract of M , then M deformation retracts onto A and thus $A \simeq_h M$ [2, Thm. 4].

In this note we focus on homotopy equivalence, a similar but weaker notion is that of *shape equivalence*, understood as being for Čech (co)homology what homotopy theory is for singular (co)homology. Indeed, only for sufficiently “nice” spaces, singular cohomology and Čech cohomology agree. For a concise introduction to shape theory, in the sense of Fox [33], we refer the reader to [20, Sec. 3]. The crux is to work with open neighbourhoods of a set and not solely with the set itself. For instance, the Warsaw circle \mathbb{W}^1 , as studied below in Example III.8 is not homotopy equivalent to the circle \mathbb{S}^1 but the two spaces are shape equivalent.

III. COFIBRATIONS

It follows from Theorem II.5 that for A to be homotopy equivalent to $\mathcal{D}_\Sigma(A)$, it suffices for A to be a neighbourhood deformation retract of M . We will appeal to *cofibrations* to capture this property.

The theory of cofibrations emerges from the so-called “*extension problem*”, that is, understanding when a continuous map $f : A \subseteq X \rightarrow Y$ can be extended from A to all of X . A typical counterexample is any map $f : \partial\mathbb{D}^n \rightarrow Y$ of nonzero degree, that is, when $\deg(f) \neq 0$, f cannot be extended from the n -sphere $\mathbb{S}^n \simeq \partial\mathbb{D}^{n+1}$ to the $(n+1)$ -disk \mathbb{D}^{n+1} [29, Sec. 2.4].

Now, cofibrations tell us, loosely speaking, if maps that can be extended, lead to homotopies that be extended. To define this, we need the following. Let X be a topological space and $A \subseteq X$, then, a pair (X, A) has the *homotopy extension property* (HEP) when, for any Y , the diagram

$$\begin{array}{ccc} (A \times [0, 1]) \cup (X \times \{0\}) & \longrightarrow & Y \\ \downarrow & \nearrow & \\ X \times [0, 1] & & \end{array} \quad (1)$$

can always be completed (the *dotted arrow* $\cdots \rightarrow$ can be found) to be commutative. Differently put, given a homotopy $H : A \times [0, 1] \rightarrow Y$ and some map $g : X \rightarrow Y$ such that $H(\cdot, 0) = g|_A$, one needs to be able to extend the homotopy from A to X . Pick $Y = (A \times [0, 1]) \cup (X \times \{0\})$, then we see that (X, A) having the HEP implies that $(A \times [0, 1]) \cup (X \times \{0\})$ is a retract of $X \times [0, 1]$. On the other hand, one

can show that the existence of such a retract implies that (X, A) has the HEP, that is, these two notions are equivalent, see Theorem III.2 below. See also [34, p. 13] for a stronger result.

Then, a continuous map $i : A \rightarrow X$ is said to be a *cofibration*² if the following commutative diagram

$$\begin{array}{ccc} A \times \{0\} & \hookrightarrow & A \times [0, 1] \\ \downarrow & & \downarrow \\ X \times \{0\} & \hookrightarrow & X \times [0, 1] \end{array} \quad \begin{array}{c} \nearrow \\ \searrow \end{array} \quad \begin{array}{c} \nearrow \\ \searrow \end{array} \quad Y \quad (2)$$

can be completed for any Y . Loosely speaking, the map i is a cofibration when it has the HEP³.

Next, we need a slight variation of the aforementioned notions of retraction, that of a *neighbourhood deformation retract pair* (NDR pair).

Definition III.1 (NDR pair). *A pair (X, A) is said to be a NDR pair if:*

- (i) *there is a continuous map $u : X \rightarrow [0, 1]$ such that $A = u^{-1}(0)$; and*
- (ii) *there is a homotopy $H : X \times [0, 1] \rightarrow X$ such that $H(\cdot, 0) = \text{id}_X$, $H(x, s) = x \ \forall x \in A, \forall s \in [0, 1]$ and $H(x, 1) \in A$ if $u(x) < 1$.*

See that if $u(x) < 1 \ \forall x \in X$, then, A is a strong deformation retract of X . In general, however, we cannot assume u to be of this form. See that for (X, A) to be a NDR pair, A must be closed. Now, a useful result is the following.

Theorem III.2 ([35, Ch. 6]). *Let A be closed in X , then, the following are equivalent:*

- (i) *the inclusion $\iota_A : A \hookrightarrow X$ is a cofibration;*
- (ii) *$(A \times [0, 1]) \cup (X \times \{0\})$ is a retract of $X \times [0, 1]$;*
- (iii) *(X, A) is a NDR pair.*

To be somewhat self-contained, we provide intuition regarding Item (ii) and Item (iii). In particular, we highlight continuity.

Proof (sketch). Suppose we have the retract $r : X \times [0, 1] \rightarrow (A \times [0, 1]) \cup (X \times \{0\})$. Define the projections $\pi_1 : X \times [0, 1] \rightarrow X$ and $\pi_2 : X \times [0, 1] \rightarrow [0, 1]$. Next, define the map $u : X \rightarrow [0, 1]$ via

$$u(x) = \sup_{\tau \in [0, 1]} \{\tau - \pi_2(r(x, \tau))\}, \quad (3)$$

²Our notion of cofibration is aligned with the so-called *Hurewicz cofibration*. We will not discuss *Serre cofibrations*, which are most easily understood through *fibrations*, a notion dual to that of cofibrations, e.g., see [35, Ch. 7].

³We focus on pairs (X, A) such that $A \subseteq X$, this inclusion is, however, not required for a cofibration to be well-defined. Nevertheless, to make sense of (1) one should work with $(A \times [0, 1]) \cup_i X$ instead $(A \times [0, 1]) \cup (X \times \{0\})$, that is, with the so-called “*mapping cylinder*” as further discussed below.

where the supremum is attained since $[0, 1]$ is compact and π_2 and r are continuous. Now define the homotopy $H : X \times [0, 1] \rightarrow X$ by $H(x, s) = \pi_1(r(x, s))$. Indeed, one can readily check that the pair (u, H) satisfies the properties required for (X, A) to be a NDR pair. Note that since the retract r is continuous, we have that u is not identically 0 when $X \setminus A \neq \emptyset$. The map u constructed through (3) is in fact continuous since $[0, 1]$ is compact and both π_2 and r are continuous, e.g., one can appeal to the simplest setting of Berge's maximum theorem [36]. \square

Note that in general, $(A \times [0, 1]) \cup (X \times \{0\})$ will be equivalent to the *mapping cylinder* under the inclusion map $\iota_A : A \hookrightarrow X$, that is, $M_{\iota_A} = ((A \times [0, 1]) \cup X) / \sim$ with $(a, 0) \sim \iota_A(a)$ for all $a \in A$, also denoted by $(A \times [0, 1]) \cup_{\iota_A} X$. Equivalence can possibly fail when the product and quotient topologies under consideration do not match.

For illustrative purposes, we end this section with the collection of a powerful result. Omitting the details, we recall that X is a *CW complex* when X can be constructed via iteratively “glueing” n -cells, being topological disks \mathbb{D}^n , along their boundary to a $(n-1)$ -dimensional CW complex, with a 0-dimensional CW complex being simply a set of discrete points. For instance, the circle \mathbb{S}^1 can be constructed from a single point and the interval. Then, a set $A \subseteq X$ is a *subcomplex* of the CW complex X when it is closed and a union of cells of X . For more on CW complices, we refer to [30, Ch. 0] and [24, Ch. 5].

Proposition III.3 (CW complices [30, Prop. 0.16]). *Let X be a CW complex and $A \subseteq X$ a subcomplex, then, the inclusion map $\iota_A : A \hookrightarrow X$ is a cofibration.*

Proposition III.3 hinges on $\iota_{\mathbb{S}^{n-1}} : \mathbb{S}^{n-1} \hookrightarrow \mathbb{D}^n$ being a cofibration, which can be shown via showing that $(\mathbb{D}^n, \mathbb{S}^{n-1})$ is a NDR pair, however, using a strategy of more general use, one can show there is a (strong deformation) retract from $\mathbb{D}^n \times [0, 1]$ onto $(\partial \mathbb{D}^n \times [0, 1]) \cup (\mathbb{D}^n \times \{0\})$ [30, p. 15], e.g., consider some point $(0, c) \in \mathbb{D}^n \times \mathbb{R}_{\geq 2}$ and project $(0, c)$ onto $(\partial \mathbb{D}^n \times [0, 1])$, then, the line between $(0, c)$ and this projection provides for the homotopy.

CW complices are fairly general, yet, properties that obstruct X admitting a CW decomposition are for instance: (i) X failing to be locally contractible [30, Prop. A4]; and (2) X failing to adhere to Whitehead's theorem cf. Example III.8.

A. Main result

Now, we have collected all ingredients to prove the following.

Lemma III.4 (\Leftarrow Cofibration). *Let $A \subseteq M$ be a compact, invariant, asymptotically stable, attractor with domain of attraction $\mathcal{D}_\Sigma(A)$. If $\iota_A : A \hookrightarrow \mathcal{D}_\Sigma(A)$ is a cofibration, then, A is a strong deformation retract of $\mathcal{D}_\Sigma(A)$.*

Proof. We know from Theorem II.5 that A is a weak deformation retract of $\mathcal{D}_\Sigma(A)$. Since $\iota_A : A \hookrightarrow \mathcal{D}_\Sigma(A)$ is a cofibration, we also know from Theorem III.2 that

$(\mathcal{D}_\Sigma(A), A)$ is a NDR pair. Then, recall Definition III.1 and recall the proof of Theorem III.2. Now, let $W := u^{-1}([0, 1]) \supset A$, which is open since $u : \mathcal{D}_\Sigma(A) \rightarrow [0, 1]$ is continuous, and consider the map $H|_{W \times [0, 1]}$. It is imperative to remark that this map does *not* provide a strong deformation retract from W onto A in general. The reason why we cannot conclude on the existence of such a map is that we cannot guarantee that throughout the homotopy we have $H(x, s) \in W$ for any $(x, s) \in W \times [0, 1]$. Indeed, we have a map $H|_{W \times [0, 1]} : W \times [0, 1] \rightarrow \mathcal{D}_\Sigma(A) \supseteq W$, the codomain cannot be assumed to be W . Precisely this detail was already known to Strøm cf. [34, Thm. 2], see also [37, p. 432]. Nevertheless, since A is a weak deformation retract of $\mathcal{D}_\Sigma(A)$ we know that W contains a set $V \supseteq A$ such that $\mathcal{D}_\Sigma(A)$ strongly deformation retracts onto V , that is, there is map $\tilde{H} : \mathcal{D}_\Sigma(A) \times [0, 1] \rightarrow \mathcal{D}_\Sigma(A)$ such that $\tilde{H}(x, 0) = x \ \forall x \in \mathcal{D}_\Sigma(A)$, $\tilde{H}(x, 1) \in V \ \forall x \in \mathcal{D}_\Sigma(A)$ and $\tilde{H}(x, s) = x \ \forall (x, s) \in V \times [0, 1]$. Hence, the continuous map $\tilde{H} : \mathcal{D}_\Sigma(A) \times [0, 1] \rightarrow \mathcal{D}_\Sigma(A)$ defined by

$$\tilde{H}(x, s) = \begin{cases} \tilde{H}(x, 2s) & s \in [0, \frac{1}{2}] \\ H(\tilde{H}(x, 1), 2s - 1) & s \in (\frac{1}{2}, 1] \end{cases}$$

is a homotopy and provides for the strong deformation retract of $\mathcal{D}_\Sigma(A)$ onto A . \square

To continue, we need a converse result.

Lemma III.5 (\Rightarrow Cofibration). *Suppose that M is a locally compact Hausdorff space, that Σ satisfies Assumption II.1 and let $A \subseteq M$ be a compact, invariant, asymptotically stable, attractor with domain of attraction $\mathcal{D}_\Sigma(A)$. If A is a strong deformation retract of $\mathcal{D}_\Sigma(A)$, then, $\iota_A : A \hookrightarrow \mathcal{D}_\Sigma(A)$ is a cofibration.*

Proof. We will appeal to the characterization of a cofibration as given by Theorem III.2. As A is a strong deformation retract of $\mathcal{D}_\Sigma(A)$ by assumption, then, to conclude on $(\mathcal{D}_\Sigma(A), A)$ being a NDR pair, we need to construct the map $u : \mathcal{D}_\Sigma(A) \rightarrow [0, 1]$. As A is a compact, invariant, asymptotically stable attractor, M is a locally compact Hausdorff space and Σ satisfies Assumption II.1, there is a Lyapunov function of precisely this form [26, Thm. 10.6]. \square

Theorem III.6 (Cofibrations). *Suppose that M is a locally compact Hausdorff space, that Σ satisfies Assumption II.1 and let $A \subseteq M$ be a compact, invariant, asymptotically stable, attractor with domain of attraction $\mathcal{D}_\Sigma(A)$. Then, A is a strong deformation retract of $\mathcal{D}_\Sigma(A)$ if and only if the inclusion $\iota_A : A \hookrightarrow \mathcal{D}_\Sigma(A)$ is a cofibration,*

Proof. The results follow directly by combining Lemma III.4 and Lemma III.5. \square

B. Examples

Regarding ramifications of Theorem III.6, we start with a sanity check. We know that for a linear ODE $\dot{x} = Fx$ with $F \in \mathbb{R}^{n \times n}$ a Hurwitz matrix, $A = \{0\}$ and $\mathcal{D}_\Sigma(A) = \mathbb{R}^n$. Hence, we remark that: (i) $\iota_{\{0\}} : \{0\} \hookrightarrow \mathbb{R}^n$ is a cofibration, e.g., since $(\mathbb{R}^n, \{0\})$ is a NDR pair under

the map $x \mapsto u(x) := 1 - e^{-\langle x, x \rangle}$; and (ii) \mathbb{R}^n strongly deformation retracts onto $0 \in \mathbb{R}^n$ via the map $\mathbb{R}^n \times [0, 1] \ni (x, s) \mapsto H(x, s) := s \cdot x$.

Cofibrations that are not strong deformation retracts are abundant. We start with a well-known example.

Example III.7 (Spheres and disks). *It can be shown that the inclusion $\iota_{\mathbb{S}^n} : \mathbb{S}^n \hookrightarrow \mathbb{D}^{n+1}$ is a cofibration, e.g., see the remark on CW complices above. However, \mathbb{D}^{n+1} cannot strongly deformation retract onto \mathbb{S}^n since \mathbb{S}^n and \mathbb{D}^{n+1} are not even homotopy equivalent, e.g., $\chi(\mathbb{S}^n) \neq \chi(\mathbb{D}^{n+1})$. Hence, \mathbb{S}^n cannot be a global, asymptotically stable, attractor under any semi-dynamical system $\Sigma = (\mathbb{D}^{n+1}, \varphi, \mathbb{R}_{\geq 0})$.*

We proceed with an example where we obtain topological insights through dynamical systems knowledge.

Example III.8 (The Warsaw circle). *Let $\mathbb{W}^1 := \{(0, x_2) \in \mathbb{R}^2 : x_2 \in [-1, 1]\} \cup \{(x_1, \sin(x_1^{-1})) \in \mathbb{R}^2 : x_1 \in (0, \pi^{-1})\} \cup \{\text{arc from } (0, -1) \text{ to } (\pi^{-1}, 0)\}$ denote the so-called “Warsaw circle”. The set \mathbb{W}^1 is compact, but not a manifold since \mathbb{W}^1 is not locally connected. Hastings showed⁴ that \mathbb{W}^1 can be rendered a compact, invariant, locally asymptotically stable attractor with an annular neighbourhood $A \subset \mathbb{R}^2$ as $\mathcal{D}_\Sigma(\mathbb{W}^1)$ [38]. Although the circle $\mathbb{S}^1 \subset \mathbb{R}^2$ and $\mathbb{W}^1 \subset \mathbb{R}^2$ are shape equivalent, they are not homotopy equivalent since $\pi_1(\mathbb{W}^1) \simeq 0$ while $\pi_1(\mathbb{S}^1) \simeq \mathbb{Z}$ and the fundamental group $\pi_1(\cdot)$ is homotopy invariant [24, Thm. 7.40]. As such, \mathbb{W}^1 cannot be a strong deformation retract of any annulus $A \subset \mathbb{R}^2$ it embeds in. Then, according to Theorem III.6, $\iota_{\mathbb{W}^1} : \mathbb{W}^1 \hookrightarrow A$ cannot be a cofibration.*

We recall that for inclusion maps $\iota_A : A \hookrightarrow X$ to not be a cofibration, the pair (X, A) cannot be too regular, e.g., by Proposition III.3 (X, A) cannot be a CW pair. Indeed, by the Whitehead theorem [30, Thm. 4.5], The Warsaw circle \mathbb{W}^1 is not homotopy equivalent to a CW complex. This could also be concluded by observing that CW complices must be locally path-connected.

Next, we provide an example inspired by an example from [39, p. 78–79]. Here we gain dynamical insights via topological knowledge. Before doing so, we recall the difference between the *box* and *product* topology. Let X_α be a topological space indexed by $\alpha \in A$, then, if we endow a topological space of the form $X = \prod_{\alpha \in A} X_\alpha$ with the product topology, open sets are of the form $\prod_{\alpha \in A} U_\alpha$ with U_α open in X_α and all but finitely many $U_\alpha = X_\alpha$. The box topology, on the other hand, does not require the last constraint to hold and open sets are simply of the form $\prod_{\alpha \in A} U_\alpha$ with U_α open in X_α . When A is finite, these topologies are equivalent, however, the box topology is finer than the product topology in general.

Example III.9 (The Tychonoff cube). *Let $\Omega > \aleph_0$, then, define the Tychonoff cube as $[0, 1]^\Omega$, that is, as a uncountably infinite product of the unit interval. Here we endow $[0, 1]$ with the standard topology and $[0, 1]^\Omega$ with the product topology. As such, $[0, 1]^\Omega$ is a compact Hausdorff space by Tychonoff’s*

theorem [25, Thm. 37.3] and the fact that any product of Hausdorff spaces is Hausdorff [25, Thm. 19.4]. Exploiting compactness, $\{0\}^\Omega \in [0, 1]^\Omega$ can be shown to be a strong deformation retract of $[0, 1]^\Omega$. Importantly, one cannot simply use the map $[0, 1]^\Omega \times [0, 1] \ni (x, s) \mapsto H(x, s) := s \cdot x$, which would be continuous in the box topology, but not in the product topology. Despite the strong deformation retraction, $\iota_0 : \{0\}^\Omega \hookrightarrow [0, 1]^\Omega$ is not a cofibration since otherwise, by Definition III.1 and Theorem III.2, there must be a continuous map $u : [0, 1]^\Omega \rightarrow [0, 1]$ such that $\{0\}^\Omega = u^{-1}(0)$. However, it can be shown that such a map fails to exist due to Ω being uncountable [39, p. 78–79]. Hence, Theorem III.6 implies that $\{0\}^\Omega \in [0, 1]^\Omega$ cannot be an asymptotically stable attractor, for any continuous—with respect to the product topology on $[0, 1]^\Omega$ —semi-dynamical system $\Sigma = ([0, 1]^\Omega, \varphi, \mathbb{R}_{\geq 0})$. Note that if Ω would be finite, then, the map u does exist and can be chosen to be $u : (x_1, \dots, x_\Omega) \mapsto \max_{i=1, \dots, \Omega} \{x_i\}$.

Note that Example III.9 is essentially saying that despite seemingly convenient properties of $\Sigma = ([0, 1]^\Omega, \varphi, \mathbb{R}_{\geq 0})$, a Lyapunov function fails to exist for $\{0\}^\Omega \in [0, 1]^\Omega$. Concurrently, this example shows that even a strong notion of homotopy equivalence can be insufficient to conclude on the existence of an asymptotically stable attractor.

Although, in general, a metrizable space must be merely countably locally finite (σ -locally finite) [25, Thm. 40.3], compact metric spaces must be second countable. Hence, $[0, 1]^{\Omega > \aleph_0}$ is not metrizable since $\Omega > \aleph_0$ obstructs second countability. Similarly, one can consider the topology of pointwise convergence. Regardless, Example III.9 illustrates where to look for counterexamples. Indeed, as $[0, 1]^{\Omega > \aleph_0}$ is not a normed space and in particular not a Hilbert space, it does not fit into common analysis frameworks, e.g., [40].

It turns out that Theorem III.6 covers known results in case A is an embedded submanifold of M . We will assume all our manifolds under consideration to be C^∞ -smooth and second countable. In that case, let $A \subseteq M$ be a closed embedded submanifold, then, one appeals to the existence of a *tubular neighbourhood* [27, Thm. 6.24] to show that $(\mathcal{D}_\Sigma(A), A)$ compromises a NDR pair. Hence, using the following proposition, [2, Prop. 10] follows as a corollary to Theorem III.6, see also [32].

Proposition III.10 (Submanifolds). *Let A be a compact embedded submanifold of M , then, $\iota_A : A \hookrightarrow M$ is a cofibration.*

Our last example pertains to compositions, indicating that Theorem III.6 can be applied to subsystems.

Example III.11 (Compositions). *Cofibrations are closed under composition. Let $i_1 : A \rightarrow B$ and $i_2 : B \rightarrow C$ be cofibrations, then, $i := i_2 \circ i_1 : A \rightarrow C$ is a cofibration. To*

⁴Although a substantial part of the proof is left to the reader.

see this, consider the diagram

$$\begin{array}{ccc}
 A \times \{0\} & \xrightarrow{\quad} & A \times [0, 1] \\
 \downarrow & & \downarrow \\
 B \times \{0\} & \xrightarrow{\quad} & B \times [0, 1] \\
 \downarrow & & \downarrow \\
 C \times \{0\} & \xrightarrow{\quad} & C \times [0, 1]
 \end{array}
 \quad (4)$$

α_1
 β_1
 α_2
 β_2

For any triple (Y, α_1, α_2) there is some appropriate map β_1 completing (2) for the cofibration $i_1 : A \rightarrow B$, but for any triple (Y, β_1, β_2) the diagram (4) can be completed since $i_2 : B \rightarrow C$ is a cofibration.

IV. FUTURE WORK

Several directions of future work are: (i) discrete systems of the form $\Sigma = (M, \varphi, \mathbb{Z}_{\geq 0})$; (ii) exploiting duality (fibrations); (iii) extensions to other notions of stability; (iv) developing computational tool (homology); (v) relaxing the invariance assumption, which is possible; and (vi) addressing stochastic systems in a meaningful way.

A. Closed attractors

Several results hold when the compactness assumption on $A \subseteq M$ is relaxed to A being merely *closed*, e.g., see [26], [41], this generalization is future work. We do emphasize that the generalization is not trivial. Consider for instance [32, Ex. 22] with $M = \mathbb{R}^2 \setminus \{(1, 0)\}$ and $A = \mathbb{S}^1 \setminus \{(1, 0)\} \subset M$. There, the authors construct a vector field on M such that A is a global asymptotically stable attractor, with $\mathcal{D}_\Sigma(A) = M \setminus \{(0, 0)\}$. So, although A is an attractor and $\iota_A : A \hookrightarrow \mathcal{D}_\Sigma(A)$ is a cofibration due to Proposition III.10, A cannot be a strong deformation retract of $\mathcal{D}_\Sigma(A)$ since those sets are not homotopy equivalent. Indeed, A is *not* compact and one cannot simply appeal to Theorem II.5. The intuition is that for attractors of this form, limits need not be attained and as such stability does not provide for a homotopy between A and $\mathcal{D}_\Sigma(A)$. Formally speaking, the proof of Theorem II.5 exploits compactness of the sublevel sets of the corresponding Lyapunov function and implicitly *Cantor's intersection theorem*, which fails to be generally true for closed sets. See also [21, Counterex. 1].

REFERENCES

- [1] J. Dydak, "Ideas and influence of Karol Borsuk," *Wiadom. Mat.*, vol. 48, no. 2, pp. 81–95, 2012.
- [2] E. Moulay and S. P. Bhat, "Topological properties of asymptotically stable sets," *Nonlinear Anal.-Theor.*, vol. 73, no. 4, pp. 1093–1097, 2010.
- [3] E. D. Sontag, *Mathematical control theory: deterministic finite dimensional systems*. New York: Springer, 1998.
- [4] S. P. Bhat and D. S. Bernstein, "A topological obstruction to continuous global stabilization of rotational motion and the unwinding phenomenon," *Syst. Control Lett.*, vol. 39, pp. 63–70, 2000.
- [5] N. P. Bhatia, "On asymptotic stability in dynamical systems," *Math. Syst. Theory*, vol. 1, no. 2, pp. 113–127, 1967.
- [6] G. D. Birkhoff, *Dynamical systems*. Providence: American Mathematical Society, 1927.

- [7] V. Nemytskii and V. Stepanov, *Qualitative theory of differential equations*. Princeton: Princeton University Press, 1960.
- [8] J. Auslander, N. Bhatia, and P. Siebert, "Attractors in dynamical systems," *Bol. Soc. Mat. Mex.*, vol. 9, pp. 55–66, 1964.
- [9] F. W. Wilson Jr, "The structure of the level surfaces of a Lyapunov function," *J. Differ. Equ.*, vol. 3, no. 3, pp. 323–329, 1967.
- [10] W. Hahn, *Stability of motion*. Berlin: Springer, 1967.
- [11] N. P. Bhatia and G. P. Szegő, *Stability theory of dynamical systems*. Berlin: Springer, 1970.
- [12] C. C. Conley, *Isolated invariant sets and the Morse index*. Providence: American Mathematical Society, 1978.
- [13] J. Milnor, "On the concept of attractor," *Comm. Math. Phys.*, vol. 99, pp. 177–195, 1985.
- [14] W. Jongeneel and E. Moulay, *Topological Obstructions to Stability and Stabilization: History, Recent Advances and Open Problems*. Cham: Springer Nature, 2023.
- [15] B. Garay, "Strong cellularity and global asymptotic stability," *Fund. Math.*, vol. 2, no. 138, pp. 147–154, 1991.
- [16] A. Giraldo, M. A. Morón, F. R. Del Portal, and J. Sanjurjo, "Some duality properties of non-saddle sets," *Topol. Appl.*, vol. 113, no. 1–3, pp. 51–59, 2001.
- [17] A. Giraldo and J. Sanjurjo, "Singular continuations of attractors," *SIAM J. Appl. Dyn.*, vol. 8, no. 2, pp. 554–575, 2009.
- [18] M. D. Kvalheim and D. E. Koditschek, "Necessary conditions for feedback stabilization and safety," *J. Geom. Mech.*, vol. 14, no. 4, pp. 659–693, 2022.
- [19] B. Günther and J. Segal, "Every attractor of a flow on a manifold has the shape of a finite polyhedron," *Proc. Am. Math. Soc.*, vol. 119, no. 1, pp. 321–329, 1993.
- [20] L. Kapitanski and I. Rodnianski, "Shape and Morse theory of attractors," *Commun. Pure Appl. Math.*, vol. 53, no. 2, pp. 218–242, 2000.
- [21] W. Yao, B. Lin, B. D. Anderson, and M. Cao, "The domain of attraction of the desired path in vector-field guided path following," *IEEE Trans. Automat. Contr.*, 2023.
- [22] C. G. Mayhew and A. R. Teel, "On the topological structure of attraction basins for differential inclusions," *Syst. Control Lett.*, vol. 60, no. 12, pp. 1045–1050, 2011.
- [23] D. Li and A. Qi, "Morse equation of attractors for nonsmooth dynamical systems," *J. Differ. Equ.*, vol. 253, no. 11, pp. 3081–3100, 2012.
- [24] J. M. Lee, *Introduction to Topological Manifolds*. New York: Springer, 2011.
- [25] J. Munkres, *Topology*. Essex: Pearson Education Limited, 2014.
- [26] N. P. Bhatia and O. Hájek, *Local semi-dynamical systems*. Berlin: Springer, 2006.
- [27] J. M. Lee, *Introduction to Smooth Manifolds*. New York: Springer, 2012.
- [28] M. Hurley, "Attractors: Persistence, and density of their basins," *T. Am. Math. Soc.*, vol. 269, no. 1, pp. 247–271, 1982.
- [29] V. Guillemin and A. Pollack, *Differential Topology*. Providence: American Mathematical Society, 2010.
- [30] A. Hatcher, *Algebraic Topology*. Cambridge: Cambridge University Press, 2002.
- [31] S. T. Hu, *Theory of retracts*. Detroit: Wayne State University Press, 1965.
- [32] B. Lin, W. Yao, and M. Cao, "On Wilson's theorem about domains of attraction and tubular neighborhoods," *Syst. Control Lett.*, vol. 167, p. 105322, 2022.
- [33] F. R. H., "On shape," *Fund. Math.*, vol. 74, pp. 47–71, 1972.
- [34] A. Strøm, "Note on cofibrations," *Math. Scand.*, vol. 19, no. 1, pp. 11–14, 1966.
- [35] J. P. May, *A concise course in algebraic topology*. Chicago: University of Chicago Press, 1999.
- [36] C. Berge, *Topological spaces*. Oliver & Boyd, 1963.
- [37] G. E. Bredon, *Topology and geometry*. New York: Springer, 1993.
- [38] H. Hastings, "A higher-dimensional Poincaré-Bendixson theorem," *Glas. Mat.*, vol. 14, no. 34, pp. 263–268, 1979.
- [39] T. tom Dieck, K. H. Kamps, and D. Puppe, *Homotopietheorie*. Berlin: Springer-Verlag, 1970.
- [40] R. F. Curtain and H. Zwart, *An introduction to infinite-dimensional linear systems theory*. New York: Springer, 2012.
- [41] Y. Lin, E. D. Sontag, and Y. Wang, "A smooth converse Lyapunov theorem for robust stability," *SIAM J. Control Optim.*, vol. 34, no. 1, pp. 124–160, 1996.